

tivdak<sup>®</sup>  
tisotumab vedotin-tftv  
for injection 40 mg



# Tivdak Eye Drop Tracker

When receiving Tivdak, it's important to take care of your eyes. This resource will help you keep track of your eye drop schedule.

## What is TIVDAK™?

TIVDAK is a prescription medicine used to treat adults with cervical cancer:

- that has returned or has spread to other parts of the body, **and**
- who have received chemotherapy that did not work or is no longer working

TIVDAK was FDA-approved based on a clinical study that measured how many patients had a tumor response and how long that response lasted. There is another study with TIVDAK to confirm the clinical benefit.

## Select Important Safety Information

### What is the most important information I should know about TIVDAK?

**Eye problems** are common with TIVDAK and can be serious. TIVDAK can cause changes to the surface of your eye that can lead to dry eyes, eye redness, eye irritation, corneal ulcers, blurred vision, and severe vision loss. **Tell your healthcare provider if you develop new or worsening vision changes or eye problems during treatment.**

Continue reading information about eye problems throughout brochure.

Please see additional [Important Safety Information](#) on pages 2-3 and [Important Facts](#) about TIVDAK, including IMPORTANT WARNING at the end of this document.

## Important Safety Information (continued)

Your healthcare provider will send you to an eye specialist to check your eyes before you start treatment with TIVDAK, before each dose of TIVDAK, and as needed for any new or worsening signs and symptoms of eye problems.

Your healthcare provider will prescribe 3 different types of eye drops before you start treatment with TIVDAK. **Bring the eye drops with you to each infusion and use them as directed by your healthcare provider to reduce your risk of eye problems:**

- You should use steroid eye drops before each infusion and as prescribed for 72 hours after each infusion.
- You should use vasoconstrictor eye drops right before each infusion.
- You should use lubricating eye drops throughout treatment and for 30 days after your last dose of TIVDAK.

Do not wear contact lenses throughout your treatment with TIVDAK unless you are told to use them by your eye specialist.

### What are the possible side effects of TIVDAK?

**Serious side effects of TIVDAK may include:**

**Peripheral neuropathy (nerve problems)** can occur with TIVDAK and can be serious. Tell your healthcare provider right away if you have new or worsening numbness or tingling in your hands or feet or muscle weakness.

**Bleeding (hemorrhage)** can occur with TIVDAK and can be serious. Tell your healthcare provider right away if you get signs or symptoms of bleeding during treatment with TIVDAK including blood in your stool or black stool (looks like tar), blood in your urine, cough up or vomit blood, unusual vaginal bleeding, any unusual or heavy bleeding.

**Lung problems.** TIVDAK may cause severe or life-threatening inflammation of the lungs that can lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, or cough.

**The most common side effects of TIVDAK include:**

- Decreased red blood cell and white blood cell counts
- Tiredness
- Nausea
- Numbness or tingling in your hands or feet, or muscle weakness (peripheral neuropathy)
- Hair loss (alopecia)
- Nosebleed (epistaxis)
- Eye problems (conjunctival disorders)
- Bleeding (hemorrhage)
- Changes in kidney function blood tests
- Dry eye
- Abnormal blood clotting test results
- Diarrhea
- Rash

Your healthcare provider may decrease your dose of TIVDAK, temporarily stop, or completely stop treatment if you have side effects.

**Important Safety Information continued on next page.**

## Important Safety Information (continued)

### What should I tell my healthcare provider before receiving TIVDAK?

Tell your healthcare provider about all your medical conditions, including if you:

- have a history of vision or eye problems
- have numbness or tingling in your hands or feet
- have bleeding problems
- have liver problems
- are pregnant or plan to become pregnant, TIVDAK can harm your unborn baby
- are breastfeeding (nursing) or plan to breastfeed. It is not known if TIVDAK passes into your breast milk. Do not breastfeed during treatment and for at least 3 weeks after the last dose of TIVDAK

#### Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with TIVDAK.
- You should use an effective method of birth control during your treatment and for at least 2 months after the last dose of TIVDAK.

#### Males with female partners who are able to become pregnant:

- You should use effective birth control during treatment and for 4 months after the last dose of TIVDAK

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

These are not all the possible side effects of TIVDAK. Discuss side effects with your healthcare provider. You are encouraged to report side effects of prescription drugs to the FDA. Visit [www.fda.gov/Safety/MedWatch](http://www.fda.gov/Safety/MedWatch) or call 1-800-FDA-1088.

Please see [Important Facts](#) about TIVDAK including **IMPORTANT WARNING** at the end of this document.

**tivdak**<sup>®</sup>   
tisotumab vedotin-tftv  
for injection 40 mg

# Tivdak Eye Drop Tracker

Your doctor will prescribe 3 different types of topical eye drops.



Vasoconstrictor  
Drops



Steroid  
Drops



Lubricating  
Drops

This calendar will help you keep track of your eye drop schedule. See instructions below:

**1** Fill in the date for each day.

**2** 3 vasoconstrictor drops (or as prescribed) will be applied once, right before your infusion, by your care team.

**4** Continue to apply steroid eye drops 1 drop per eye, 3x a day or as instructed by your doctor, for Day 2 and 3 after each infusion.

INFUSION CYCLE	Day 1	Day 2	Day 3
<b>Date</b>	INFUSION DAY <i>6/21</i>	<i>6/22</i>	<i>6/23</i>
<b>Vasoconstrictor Drops</b>		—	—
<b>Steroid Drops</b>			
<b>Lubricating Drops</b>	Remember to take your lubricating drops for the duration of your treatment until 30 days after your last dose of Tivdak.		
<b>Notes:</b> <i>Add notes here</i>			

**3** Take your first steroid drop in each eye at your infusion appointment. Your care team may do this for you.

**5** Take notes on how you're feeling.

Your care team will decide how many treatments you will need

Please see [Important Safety Information](#) on pages 2-3 and [Important Facts](#) about TIVDAK, including IMPORTANT WARNING at the end of this document.

Mark each circle with an X once the drops are applied.  
Always take your eye drops as directed by your doctor

INFUSION CYCLE	Day 1	Day 2	Day 3
Date	INFUSION DAY		
Vasoconstrictor Drops	<input type="checkbox"/>	—	—
Steroid Drops	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
Lubricating Drops	Remember to take your lubricating drops for the duration of your treatment until 30 days after your last dose of Tivdak. →		
Notes:			

INFUSION CYCLE	Day 1	Day 2	Day 3
Date	INFUSION DAY		
Vasoconstrictor Drops	<input type="checkbox"/>	—	—
Steroid Drops	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
Lubricating Drops	Remember to take your lubricating drops for the duration of your treatment until 30 days after your last dose of Tivdak. →		
Notes:			

INFUSION CYCLE	Day 1	Day 2	Day 3
Date	INFUSION DAY		
Vasoconstrictor Drops	<input type="checkbox"/>	—	—
Steroid Drops	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	<input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>
Lubricating Drops	Remember to take your lubricating drops for the duration of your treatment until 30 days after your last dose of Tivdak. →		
Notes:			

Please see [Important Safety Information](#) on pages 2-3 and [Important Facts](#) about TIVDAK, including IMPORTANT WARNING at the end of this document.

# Reminders

- Always take your eye drops as directed by your doctor
- Take your lubricating drops for the duration of treatment and for 30 days after your last dose of Tivdak
- Monitor your eyes every day—tell your care team right away if you notice any new or worsening eye problems, such as swelling, redness, itchy eyes, eye pain, blurry vision, watery eyes, dry eyes, light sensitivity, or severe vision loss
- Print out more Tivdak Eye Drop Trackers at [tivdak.com](http://tivdak.com)
- Bring your Tivdak Eye Drop Tracker and all of your eye drops to your infusion appointments
- Keep track of your eye drops and refill prescriptions when needed
- Always ask your care team any questions you have throughout your treatment with Tivdak. Your care team is there to help

---

To learn more visit [tivdak.com](http://tivdak.com)

---

Tivdak and its logo are trademarks of Seagen Inc. Seagen and its logo are US registered trademarks of Seagen Inc. Genmab; the Y-shaped Genmab logo; and Genmab in combination with the Y-shaped Genmab logo are trademarks for Genmab A/S and/or its subsidiaries.

©2021 Seagen Inc. and Genmab US, Inc. All rights reserved. US-TVP-21-144-MT

# Important Facts

This is only a summary of important information about TIVDAK. Talk to your care team or pharmacist to learn more.

**tivdak**<sup>™</sup>  
tisotumab vedotin-tftv  
for injection 40 mg

## What is the most important information I should know about TIVDAK?

TIVDAK can cause serious side effects, including:

**Eye problems.** Eye problems are common with TIVDAK, and can also be serious. TIVDAK can cause changes to the surface of your eye that can lead to dry eyes, eye redness, eye irritation, corneal ulcers, blurred vision, and severe vision loss. **Tell your healthcare provider if you develop new or worsening vision changes or eye problems during treatment with TIVDAK.**

- Your healthcare provider will send you to an eye specialist to check your eyes before you start treatment with TIVDAK, before each dose of TIVDAK, and as needed for any new or worsening signs and symptoms of eye problems.
- Your healthcare provider will prescribe 3 different types of eye drops before you start treatment with TIVDAK. **Bring the eye drops with you to each infusion and use them as directed by your healthcare provider to reduce your risk of eye problems:**
  - You should use steroid eye drops before each infusion and as prescribed for 72 hours after each infusion.
  - You should use vasoconstrictor eye drops right before each infusion.
  - You should use lubricating eye drops throughout treatment and for 30 days after your last dose of TIVDAK.
- Do not wear contact lenses throughout your treatment with TIVDAK unless you are told to use them by your eye specialist.
- See “**What are the possible side effects of TIVDAK?**” for more information about side effects.

## What is TIVDAK?

TIVDAK is a prescription medicine used to treat adults with cervical cancer:

- that has returned or has spread to other parts of the body, **and**
- who have received chemotherapy that did not work or is no longer working.

It is not known if TIVDAK is safe and effective in children.

## Before receiving TIVDAK

**Tell your healthcare provider about all of your medical conditions, including if you:**

- have a history of vision or eye problems
- have numbness or tingling in your hands or feet
- have bleeding problems
- have liver problems
- are pregnant or plan to become pregnant. TIVDAK can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TIVDAK.

### Females who are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with TIVDAK.
- You should use an effective birth control during treatment and for 2 months after your last dose of TIVDAK.

### Males with female partners who are able to become pregnant:

- You should use an effective birth control during treatment and for 4 months after your last dose of TIVDAK.
- are breastfeeding or plan to breastfeed. It is not known if TIVDAK passes into your breast milk. Do not breastfeed during treatment and for 3 weeks after your last dose of TIVDAK.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Taking TIVDAK with certain other medicines may cause side effects.

## How will I receive TIVDAK?

- TIVDAK will be given to you by intravenous (IV) infusion into your vein over 30 minutes.
- TIVDAK is usually given every 3 weeks.
- Your healthcare provider will decide how many infusions you need.
- Your healthcare provider will put cold packs on your eyes during each infusion.
- Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with TIVDAK if you have side effects.

## What are the possible side effects of TIVDAK?

TIVDAK can cause serious side effects, including:

- See “**What is the most important information I should know about TIVDAK?**”
- **Peripheral neuropathy.** Nerve problems called peripheral neuropathy are common with TIVDAK, and can also be serious. Tell your healthcare provider right away if you get numbness or tingling in your hands or feet or muscle weakness.
- **Bleeding (hemorrhage).** Bleeding problems are common with TIVDAK, and can also be serious. Tell your healthcare provider right away if you get signs or symptoms of bleeding during treatment with TIVDAK, including:
  - blood in your stools or black stools (looks like tar)
  - blood in your urine
  - cough up or vomit blood
  - unusual vaginal bleeding
  - any unusual or heavy bleeding
- **Lung problems.** TIVDAK may cause severe or life-threatening inflammation of the lungs that can lead to death. Tell your healthcare provider right away if you get new or worsening symptoms, including trouble breathing, shortness of breath, or cough.

### The most common side effects of TIVDAK include:

- decreased red blood cell and white blood cell counts
- tiredness
- nausea
- hair loss (alopecia)
- nosebleed
- changes in kidney function blood tests
- dry eye
- abnormal blood clotting test results
- diarrhea
- rash

# Important Facts

This is only a summary of important information about TIVDAK. Talk to your care team or pharmacist to learn more.

**tivdak**<sup>™</sup>  
tisotumab vedotin-tftv  
for injection 40 mg

## What are the possible side effects of TIVDAK? (continued)

TIVDAK may cause fertility problems in males, which may affect your ability to father children. Talk to your healthcare provider if this is a concern for you.

These are not all the possible side effects of TIVDAK.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

## Get more information

- This is only a summary of key information about TIVDAK
- Talk to your healthcare provider or pharmacist to learn more
- Go to [tivdak.com](http://tivdak.com) or call 1-855-4SEAGEN
- If you need help paying for your medicine, visit [SeagenSecure.com](http://SeagenSecure.com) for program information

Visit [tivdak.com](http://tivdak.com) to learn more or call 1-855-4SEAGEN.